JOURNAL OF CHROMATOGRAPHY A

VOL. 761, NOS. 1 + 2 14 FEBRUARY 1997

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

OBITUARY 1

REGULAR PAPERS

Column Liquid Chromatography

Ideal model of chromatography applied to charged solutes in reversed-phase liquid chromatography
by I. Hägglund (Södertälje, Sweden) and J. Ståhlberg (Stockholm, Sweden) (Received 7 October 1996) 3

Method for determination of the chromatographic surface area of reversed-phase surfaces
by I. Hägglund (Södertälje, Sweden) and J. Ståhlberg (Stockholm, Sweden) (Received 7 October 1996) 13

Liquid adsorption chromatography of polyethers. Theory and experiment
by B. Trathnigg and M. Kollroser (Graz, Austria) and A. Gorbunov and A. Skvortsov (St. Petersburg, Russia) (Received 16 September 1996) 21

Estimation of pore diameter for intraparticle fluid flow in bidisperse porous chromatographic particles
by G.A. Heeter and A.I. Liapis (Rolla, MO, USA) (Received 16 September 1996) 35

On the reproducibility of column performance in liquid chromatography and the role of the packing density
by B.J. Stanley and C.R. Foster (San Bernardino, CA, USA) and G. Guiochon (Knoxville and Oak Ridge, TN, USA) (Received 25 September 1996) 41

Comparison of chromatographic systems for triterpenoids from Neem (*Azadirachta indica*) seeds
by S. Johnson and E.D. Morgan (Keele, UK) (Received 16 September 1996) 53

Predictable chromatographic separations of enantiomers: aryl allenic acids and their derivatives
by W.H. Pirkle and M.E. Koscho (Urbana, IL, USA) (Received 15 August 1996) 65

Rapid determination of amphetamines by high-performance liquid chromatography with UV detection
by F. Sadeghipour (Genève, Switzerland), C. Giroud and L. Rivier (Lausanne, Switzerland) and J.-L. Veuthey (Genève, Switzerland) (Received 16 October 1996) 71

Study of retention, efficiency and selectivity in chiral ligand-exchange chromatography with a dynamically coated stationary phase
by M. Remelli, P. Fornasari and F. Pulidori (Ferrara, Italy) (Received 19 September 1996) 79

Partitioning of hydrophobic amino acids and oligopeptides in aqueous two-phase system containing self-aggregating block copolymer. Effects of temperature, salts and surfactants
by M. Svensson, F. Joabsson, P. Linse and F. Tjerneld (Lund, Sweden) (Received 26 September 1996) 91

High-performance liquid chromatographic methods for separation of enantiomers of alicyclic β-amino acids
by A. Péter, G. Török, P. Csomós, M. Péter, G. Bernáth and F. Fülöp (Szeged, Hungary) (Received 27 September 1996) 103

Reversed-phase liquid chromatographic separation of platinum(II) complexes of methionine- and histidine-containing peptides using perfluorinated carboxylic acids as ion-pairing reagents
by A.F.M. Siebert, C.D.W. Fröhling and W.S. Sheldrick (Bochum, Germany) (Received 26 September 1996) 115

Investigation into the chiral recognition mechanism of N-arylthiazolin-2(thi)one atropisomers on Chiralcel OJ by factorial design and lipophilicity approaches
by C. Roussel and C. Suteu (Marseilles, France) (Received 17 September 1996) 129

(Continued overleaf)

Contents (continued)

Reversed-phase high-performance liquid chromatography of pyrrolidinedithiocarbamate complexes of mercuric species using amperometric and coulometric detection
by M.P. da Silva, J.R. Procopio and L. Hernández (Madrid, Spain) (Received 26 September 1996) 139

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. II. Solubility effects in the analysis of polyester resins under reversed-phase conditions
by H.J.A. Philipsen (Venlo, Netherlands) and M.R. de Cooker, H.A. Claessens, B. Klumperman and A.L. German (Eindhoven, Netherlands) (Received 24 September 1996) 147

Preconcentration determination of inorganic anions and organic acids in power plant waters. Separation optimization through control of column capacity and selectivity
by M. Toofan, J.R. Stillian, C.A. Pohl and P.E. Jackson (Sunnyvale, CA, USA) (Received 4 October 1996) 163

Gas Chromatography

Helium direct current discharge ionization detector for gas chromatography
by Q. Jin, W. Yang, A. Yu, X. Tian and F. Wang (Changchun, China) (Received 17 September 1996) 169

Enantioselective capillary gas chromatography and capillary supercritical fluid chromatography on an immobilized γ-cyclodextrin derivative
by H. Grosenick and V. Schurig (Tübingen, Germany) (Received 19 September 1996) 181

Gas chromatography of Titan's atmosphere. VIII. Analysis of permanent gases with carbon molecular sieve packed capillary columns
by A. Aflalaye and R. Sternberg (Créteil, France), D. Coscia (Verrières le Buisson, France), F. Raulin (Créteil, France) and C. Vidal-Madjar (Thiais, France) (Received 17 September 1996) 195

Solid-phase microextraction of volatile organic compounds. Estimation of the sorption equilibrium from the Kováts index, effect of salinity and humic acids and the study of the kinetics by the development of an "agitated/static layer" model
by J. Dewulf, H. Van Langenhove and M. Everaert (Ghent, Belgium) (Received 25 September 1996) 205

Procedures for analysing phenolic metabolites of polychlorinated dibenzofurans, -dibenzo-*p*-dioxins and -biphenyls extracted from a microsomal assay: optimising solid-phase adsorption clean-up and derivatisation methods
by M.J.C. Rozemeijer, K. Olie and P. de Voogt (Amsterdam, Netherlands) (Received 26 September 1996) 219

Molecular analysis of the sulphur components in a light cycle oil of a catalytic cracking unit by gas chromatography with mass spectrometric and atomic emission detection
by G.A. Depauw and G.F. Froment (Ghent, Belgium) (Received 30 September 1996) 231

Electrophoresis

Chiral separation of neutral species by capillary electrophoresis. Evaluation of a theoretical model
by S. Surapaneni, K. Ruterbories and T. Lindstrom (Indianapolis, IN, USA) (Received 7 October 1996) 249

Capillary electrophoretic separation of chlorophenols using amperometric detection
by A. Hilmi, J.H.T. Luong and A.-L. Nguyen (Montreal, Canada) (Received 16 September 1996) 259

Chiral resolution of diols by capillary electrophoresis using borate–cyclodextrin complexation
by T. Jira and A. Bunke (Greifswald, Germany) and M.G. Schmid and G. Gübitz (Graz, Austria) (Received 17 September 1996) 269

Analysis of prostaglandins by micellar electrokinetic capillary chromatography
by Y.-Z. Hsieh and K.-L. Kuo (Hsinchu, Taiwan) (Received 16 September 1996) 277

Use of capillary electrophoresis and high-performance liquid chromatography for monitoring of glycosylation of the peptides dalargin and desmopressin
by J. Jiskra, V. Pacáková, M. Tichá, K. Štulík and T. Barth (Prague, Czech Republic) (Received 19 September 1996) 285

Improved separation of six harmane alkaloids by high-performance capillary electrophoresis
by J. Cheng (Philadelphia, PA, USA) and K.R. Mitchelson (Indooroopilly, Australia) (Received 5 September 1996) 297

Separation of retinoids by micellar electrokinetic capillary chromatography
by Y.-Z. Hsieh and K.-L. Kuo (Hsinchu, Taiwan) (Received 30 September 1996) 307

Contents (continued)

SHORT COMMUNICATIONS

Column Liquid Chromatography

Quantitative analysis of flavonoids by reversed-phase high-performance liquid chromatography
by A. Crozier (Glasgow, UK), E. Jensen (Tromsø, Norway) and M.E.J. Lean and M.S. McDonald (Glasgow, UK) (Received 27 August 1996) 315

Direct determination of bromacil and diuron residues in environmental water samples by coupled-column liquid chromatography and large-volume injection
by J.V. Sancho, C. Hidalgo and F. Hernández (Castelló, Spain) (Received 25 September 1996) 322

Gas Chromatography

Simplified multiresidue method for the determination of organophosphorus insecticides in olive oil
by P. Cabras, A. Angioni and M. Melis (Cagliari, Italy), E.V. Minelli (Araraquara, Brasil) and F.M. Pirisi (Cagliari, Italy) (Received 30 September 1996) 327

Determination of triazolam in a drug tablet by thermal desorption gas chromatography
by M. Hida, T. Mitsui, H. Ohtani and S. Tsuge (Nagoya, Japan) (Received 25 September 1996) 332

Supercritical Fluid Chromatography

Separation and identification of vitamin A acetate isomers by supercritical fluid chromatography–^{1}H NMR coupling
by U. Braumann, H. Händel and S. Strohschein (Tübingen, Germany), M. Spraul (Rheinstetten, Germany), G. Krack (Ludwigshafen, Germany), R. Ecker (Waldbronn, Germany) and K. Albert (Tübingen, Germany) (Received 16 July 1996) . . 336

ERRATUM

Erattum to ''Gas chromatographic-mass spectrometric study of photodegradation of carbamate pesticides''. [J. Chromatogr. A, 738 (1996) 225–231]
by M.J. Climent and M.A. Miranda (Valencia, Spain) (Received 27 September 1996) 341

BOOK REVIEW 343

AUTHOR INDEX VOL. 761 345

The table of contents of *Journal of Chromatography* is included in ESTOC – Elsevier Science Tables of Contents service – which can be accessed on the World Wide Web at the following URL addresses:
http://www.elsevier.nl/locate/estoc or http://www.elsevier.com/locate/estoc